

Recommended Childhood and Adolescent Immunization Schedule



UNITED STATES • 2006



Vaccine ▼	Age ►	Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	24 months	4–6 years	11–12 years	13–14 years	15 years	16–18 years
Hepatitis B ¹		HepB	HepB	HepB ¹	HepB			HepB Series							
Diphtheria, Tetanus, Pertussis ²			DTaP	DTaP	DTaP	DTaP				DTaP	Tdap	Tdap			
<i>Haemophilus influenzae</i> typeb ³			Hib	Hib	Hib ³	Hib									
Inactivated Poliovirus			IPV	IPV	IPV				IPV						
Measles, Mumps, Rubella ⁴						MMR				MMR	MMR				
Varicella ⁵						Varicella			Varicella						
Meningococcal ⁶								Vaccines within broken line are for selected populations		MPSV4	MCV4		MCV4		
Pneumococcal ⁷			PCV	PCV	PCV	PCV				PCV	PPV				
Influenza ⁸					Influenza (Yearly)			Influenza (Yearly)							
Hepatitis A ⁹									HepA Series						

This schedule indicates the recommended ages for routine administration of currently licensed childhood vaccines, as of December 1, 2005, for children through age 18 years. Any dose not administered at the recommended age should be administered at any subsequent visit when indicated and feasible. Indicates age groups that warrant special effort to administer those vaccines not previously administered. Additional vaccines may be licensed and recommended during the year. Licensed combination vaccines may be used whenever

any components of the combination are indicated and other components of the vaccine are not contraindicated and if approved by the Food and Drug Administration for that dose of the series. Providers should consult the respective ACIP statement for detailed recommendations. Clinically significant adverse events that follow immunization should be reported to the Vaccine Adverse Event Reporting System (VAERS). Guidance about how to obtain and complete a VAERS form is available at www.vaers.hhs.gov or by telephone, 800-822-7967.

 Range of recommended ages Catch-up immunization 11–12 year old assessment

- 1. Hepatitis B vaccine (HepB).** *AT BIRTH:* All newborns should receive monovalent HepB soon after birth and before hospital discharge. **Infants born to mothers who are HBsAg-positive** should receive HepB and 0.5 mL of hepatitis B immune globulin (HBIG) within 12 hours of birth. **Infants born to mothers whose HBsAg status is unknown** should receive HepB within 12 hours of birth. The mother should have blood drawn as soon as possible to determine her HBsAg status; if HBsAg-positive, the infant should receive HBIG as soon as possible (no later than age 1 week). **For infants born to HBsAg-negative mothers,** the birth dose can be delayed in rare circumstances but only if a physician's order to withhold the vaccine and a copy of the mother's original HBsAg-negative laboratory report are documented in the infant's medical record. **FOLLOWING THE BIRTHDOSE:** The HepB series should be completed with either monovalent HepB or a combination vaccine containing HepB. The second dose should be administered at age 1–2 months. The final dose should be administered at age ≥24 weeks. It is permissible to administer 4 doses of HepB (e.g., when combination vaccines are given after the birth dose); however, if monovalent HepB is used, a dose at age 4 months is not needed. **Infants born to HBsAg-positive mothers** should be tested for HBsAg and antibody to HBsAg after completion of the HepB series, at age 9–18 months (generally at the next well-child visit after completion of the vaccine series).

2. Diphtheria and tetanus toxoids and acellular pertussis vaccine (DTaP). The fourth dose of DTaP may be administered as early as age 12 months, provided 6 months have elapsed since the third dose and the child is unlikely to return at age 15–18 months. The final dose in the series should be given at age ≥4 years.

Tetanus and diphtheria toxoids and acellular pertussis vaccine (Tdap – adolescent preparation) is recommended at age 11–12 years for those who have completed the recommended childhood DTP/DTaP vaccination series and have not received a Td booster dose. Adolescents 13–18 years who missed the 11–12-year Td/Tdap booster dose should also receive a single dose of Tdap if they have completed the recommended childhood DTP/DTaP vaccination series. Subsequent **tetanus and diphtheria toxoids (Td)** are recommended every 10 years.

3. *Haemophilus influenzae* type b conjugate vaccine (Hib). Three Hib conjugate vaccines are licensed for infant use. If PRP-OMP (PedvaxHIB® or ComVax® [Merck]) is administered at ages 2 and 4 months, a dose at age 6 months is not required. DTaP/Hib combination products should not be used for primary immunization in infants at ages 2, 4 or 6 months but can be used as boosters after any Hib vaccine. The final dose in the series should be administered at age ≥12 months.

4. Measles, mumps, and rubella vaccine (MMR). The second dose of MMR is recommended routinely at age 4–6 years but may be administered during any visit, provided at least 4 weeks have elapsed since the first dose and both doses are administered beginning at or after age 12 months. Those who have not previously received the second dose should complete the schedule by age 11–12 years.
- 5. Varicella vaccine.** Varicella vaccine is recommended at any visit at or after age 12 months for susceptible children (i.e., those who lack a reliable history of chickenpox). Susceptible persons aged ≥13 years should receive 2 doses administered at least 4 weeks apart.

6. Meningococcal vaccine (MCV4). Meningococcal conjugate vaccine (MCV4) should be given to all children at the 11–12 year old visit as well as to unvaccinated adolescents at high school entry (15 years of age). Other adolescents who wish to decrease their risk for meningococcal disease may also be vaccinated. All college freshmen living in dormitories should also be vaccinated, preferably with MCV4, although **meningococcal polysaccharide vaccine (MPSV4)** is an acceptable alternative. Vaccination against invasive meningococcal disease is recommended for children and adolescents aged ≥2 years with terminal complement deficiencies or anatomic or functional asplenia and certain other high risk groups (see *MMWR* 2005;54 [RR-7]:1-21); use MPSV4 for children aged 2–10 years and MCV4 for older children, although MPSV4 is an acceptable alternative.

7. Pneumococcal vaccine. The heptavalent **pneumococcal conjugate vaccine (PCV)** is recommended for all children aged 2–23 months and for certain children aged 24–59 months. The final dose in the series should be given at age ≥12 months. **Pneumococcal polysaccharide vaccine (PPV)** is recommended in addition to PCV for certain high-risk groups. See *MMWR* 2000; 49(RR-9):1-35.

8. Influenza vaccine. Influenza vaccine is recommended annually for children aged ≥6 months with certain risk factors (including, but not limited to, asthma, cardiac disease, sickle cell disease, human immunodeficiency virus [HIV], diabetes, and conditions that can compromise respiratory function or handling of respiratory secretions or that can increase the risk for aspiration), healthcare workers, and other persons (including household members) in close contact with persons in groups at high risk (see *MMWR* 2005;54[RR-8]:1-55). In addition, healthy children aged 6–23 months and close contacts of healthy children aged 0–5 months are recommended to receive influenza vaccine because children in this age group are at substantially increased risk for influenza-related hospitalizations. For healthy persons aged 5–49 years, the intranasally administered, live, attenuated influenza vaccine (LAIV) is an acceptable alternative to the intramuscular trivalent inactivated influenza vaccine (TIV). See *MMWR* 2005;54(RR-8):1-55. Children receiving TIV should be administered a dosage appropriate for their age (0.25 mL if aged 6–35 months or 0.5 mL if aged ≥3 years). Children aged ≤8 years who are receiving influenza vaccine for the first time should receive 2 doses (separated by at least 4 weeks for TIV and at least 6 weeks for LAIV).

9. Hepatitis A vaccine (HepA). HepA is recommended for all children at 1 year of age (i.e., 12–23 months). The 2 doses in the series should be administered at least 6 months apart. States, counties, and communities with existing HepA vaccination programs for children 2–18 years of age are encouraged to maintain these programs. In these areas, new efforts focused on routine vaccination of 1-year-old children should enhance, not replace, ongoing programs directed at a broader population of children. HepA is also recommended for certain high risk groups (see *MMWR* 1999; 48[RR-12]:1-37).

Recommended Immunization Schedule for Children and Adolescents Who Start Late or Who Are More Than 1 Month Behind

UNITED STATES • 2006

The tables below give catch-up schedules and minimum intervals between doses for children who have delayed immunizations. There is no need to restart a vaccine series regardless of the time that has elapsed between doses. Use the chart appropriate for the child’s age.

CATCH-UP SCHEDULE FOR CHILDREN AGED 4 MONTHS THROUGH 6 YEARS					
Vaccine	Minimum Age for Dose 1	Minimum Interval Between Doses			
		Dose 1 to Dose 2	Dose 2 to Dose 3	Dose 3 to Dose 4	Dose 4 to Dose 5
Diphtheria, Tetanus, Pertussis	6 wks	4 weeks	4 weeks	6 months	6 months ¹
Inactivated Poliovirus	6 wks	4 weeks	4 weeks	4 weeks ²	
Hepatitis B ³	Birth	4 weeks	8 weeks (and 16 weeks after first dose)		
Measles, Mumps, Rubella	12 mo	4 weeks ⁴			
Varicella	12 mo				
<i>Haemophilus influenzae</i> type b ⁵	6 wks	4 weeks if first dose given at age <12 months	4 weeks ⁶ if current age <12 months	8 weeks (as final dose) This dose only necessary for children aged 12 months–5 years who received 3 doses before age 12 months	
		8 weeks (as final dose) if first dose given at age 12–14 months	8 weeks (as final dose) ⁶ if current age ≥12 months and second dose given at age <15 months		
		No further doses needed if first dose given at age ≥15 months	No further doses needed if previous dose given at age ≥15 mo		
Pneumococcal ⁷	6 wks	4 weeks if first dose given at age <12 months and current age <24 months	4 weeks if current age <12 months	8 weeks (as final dose) This dose only necessary for children aged 12 months–5 years who received 3 doses before age 12 months	
		8 weeks (as final dose) if first dose given at age ≥12 months or current age 24–59 months	8 weeks (as final dose) if current age ≥12 months		
		No further doses needed for healthy children if first dose given at age ≥24 months	No further doses needed for healthy children if previous dose given at age ≥24 months		

CATCH-UP SCHEDULE FOR CHILDREN AGED 7 YEARS THROUGH 18 YEARS			
Vaccine	Minimum Interval Between Doses		
	Dose 1 to Dose 2	Dose 2 to Dose 3	Dose 3 to Booster Dose
Tetanus, Diphtheria ⁸	4 weeks	6 months	6 months if first dose given at age <12 months and current age <11 years; otherwise 5 years
Inactivated Poliovirus ⁹	4 weeks	4 weeks	IPV ^{2,9}
Hepatitis B	4 weeks	8 weeks (and 16 weeks after first dose)	
Measles, Mumps, Rubella	4 weeks		
Varicella ¹⁰	4 weeks		

1. **DTaP.** The fifth dose is not necessary if the fourth dose was administered after the fourth birthday.

2. **IPV.** For children who received an all-IPV or all-oral poliovirus (OPV) series, a fourth dose is not necessary if third dose was administered at age ≥4 years. If both OPV and IPV were administered as part of a series, a total of 4 doses should be given, regardless of the child’s current age.

3. **HepB.** Administer the 3-dose series to all children and adolescents <19 years of age if they were not previously vaccinated.

4. **MMR.** The second dose of MMR is recommended routinely at age 4–6 years but may be administered earlier if desired.

5. **Hib.** Vaccine is not generally recommended for children aged ≥5 years.
6. **Hib.** If current age <12 months and the first 2 doses were PRP-OMP (PedvaxHIB® or ComVax® [Merck]), the third (and final) dose should be administered at age 12–15 months and at least 8 weeks after the second dose.

7. **PCV.** Vaccine is not generally recommended for children aged ≥5 years.

8. **Td.** Adolescent tetanus, diphtheria, and pertussis vaccine (Tdap) may be substituted for any dose in a primary catch-up series or as a booster if age appropriate for Tdap. A five-year interval from the last Td dose is encouraged when Tdap is used as a booster dose. See ACIP recommendations for further information.

9. **IPV.** Vaccine is not generally recommended for persons aged ≥18 years.

10. **Varicella.** Administer the 2-dose series to all susceptible adolescents aged ≥13 years.



CENTERS FOR DISEASE CONTROL AND PREVENTION